


CosmoLattice

School 2022 - Valencia

Non-linear dynamics of
Axion-inflation

Joanes Lizarraga & Ander Uriol



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]

...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:


$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

$\phi \rightarrow$ Pseudo-scalar axion field

[M. M. Anber, L. Sorbo (0908.4089)]

[J. Cook, L. Sorbo (1109.0022)]

[N. Barnaby, E. Pajer, M. Peloso (1110.3327)]



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]

...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

$\phi \rightarrow$ Pseudo-scalar axion field

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$


$$\tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

[M. M. Anber, L. Sorbo (0908.4089)]

[J. Cook, L. Sorbo (1109.0022)]

[N. Barnaby, E. Pajer, M. Peloso (1110.3327)]



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]

...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Simple derivative
Non covariant

$\phi \rightarrow$ Pseudo-scalar axion field

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$


$$\tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

[M. M. Anber, L. Sorbo (0908.4089)]

[J. Cook, L. Sorbo (1109.0022)]

[N. Barnaby, E. Pajer, M. Peloso (1110.3327)]



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]

...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Simple derivative
Non covariant

$\phi \rightarrow$ Pseudo-scalar axion field


$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

$$\tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

Generated from
external mechanism
breaks the shift
symmetry explicitly

[M. M. Anber, L. Sorbo (0908.4089)]
[J. Cook, L. Sorbo (1109.0022)]
[N. Barnaby, E. Pajer, M. Peloso (1110.3327)]



Basics of the model

[K. Freese, J. A. Frieman, A. V. Olinto (PRL 65,3233 1990)]
...

Axion-inflation + $\frac{\phi}{\Lambda} F \tilde{F}$

Shift symmetry $\phi \rightarrow \phi + c$

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Simple derivative
Non covariant

$\phi \rightarrow$ Pseudo-scalar axion field

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

$$\tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$


$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

Generated from
external mechanism
breaks the shift
symmetry explicitly

Interaction only through this vertex
Axion coupling!

$1/\Lambda$ Coupling constant $[m_{\text{pl}}^{-1}]$


[M. M. Anber, L. Sorbo (0908.4089)]
[J. Cook, L. Sorbo (1109.0022)]
[N. Barnaby, E. Pajer, M. Peloso (1110.3327)]



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$


Vectorial form

$$S = \int d^4x \left[\frac{1}{2} a^3 \pi_\phi^2 - \frac{1}{2} a \left(\vec{\nabla} \phi \right)^2 - \frac{1}{2} a^3 m^2 \phi^2 + \frac{1}{2} a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

@ FLRW:

$$ds^2 = -dt^2 + a^2 d\vec{x}^2$$

In cosmic time



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$


Vectorial form

$$S = \int d^4x \left[\frac{1}{2} a^3 \pi_\phi^2 - \frac{1}{2} a \left(\vec{\nabla} \phi \right)^2 - \frac{1}{2} a^3 m^2 \phi^2 + \frac{1}{2} a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

@ FLRW:

$$ds^2 = -dt^2 + a^2 d\vec{x}^2$$

In cosmic time



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Vectorial form


$$S = \int d^4x \left[\frac{1}{2} a^3 \pi_\phi^2 - \frac{1}{2} a \left(\vec{\nabla} \phi \right)^2 - \frac{1}{2} a^3 m^2 \phi^2 + \frac{1}{2} a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

@ FLRW:

$$ds^2 = -dt^2 + a^2 d\vec{x}^2$$

In cosmic time

3 vertex



Basics of the model

Action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} m_{\text{pl}}^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\phi}{4\Lambda} F_{\mu\nu} \tilde{F}^{\mu\nu} \right)$$

Vectorial form

$$S = \int d^4x \left[\frac{1}{2} a^3 \pi_\phi^2 - \frac{1}{2} a \left(\vec{\nabla} \phi \right)^2 - \frac{1}{2} a^3 m^2 \phi^2 + \frac{1}{2} a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Our choice of
the potential,
also:

$$V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3}}\phi} \right)^2$$

$$V(\phi) = V_0 \left(1 - \left(\frac{\phi}{\eta} \right)^4 \right)^2$$


$$V(\phi) = \dots$$

3 vertex

@ FLRW:

$$ds^2 = -dt^2 + a^2 d\vec{x}^2$$

In cosmic time



Basics of the model

Continuum equations of motion:


Temporal gauge

$$A_0 = 0$$

CL momentum variables

$$\tilde{\pi}_\phi = a^3 \pi_\phi, \quad \tilde{\vec{E}} = a \vec{E}$$

No Hubble friction
term in the EoMs



Basics of the model

Continuum equations of motion:

Dynamical equations:

Temporal gauge

$$A_0 = 0$$


CL momentum variables

$$\tilde{\pi}_\phi = a^3 \pi_\phi, \quad \tilde{\vec{E}} = a \vec{E}$$

No Hubble friction term in the EoMs

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$



Basics of the model

Continuum equations of motion:

Dynamical equations:

Temporal gauge

$$A_0 = 0$$

CL momentum variables

$$\tilde{\pi}_\phi = a^3 \pi_\phi, \quad \tilde{\vec{E}} = a \vec{E}$$


No Hubble friction term in the EoMs

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \vec{\nabla} \phi \cdot \vec{B} = 0$$



Basics of the model

Continuum equations of motion:

Dynamical equations:

Temporal gauge

$$A_0 = 0$$

CL momentum variables

$$\tilde{\pi}_\phi = a^3 \pi_\phi, \quad \tilde{\vec{E}} = a \vec{E}$$

No Hubble friction term in the EoMs


$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \vec{\nabla} \phi \cdot \vec{B} = 0$$

Non-linear terms



Basics of the model: linear regime

Let's go now deep inside inflation


Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda}\vec{\nabla}\phi \cdot \vec{B} = 0$$



Basics of the model: linear regime

Let's go now deep inside inflation

Dynamical equations:

Inflaton in slow-roll,
Homogeneous

+


dominated by the potential

$$\dot{\tilde{\pi}}_\phi = a \cancel{\vec{\nabla}^2} \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \cancel{\vec{\nabla} \phi} \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \cancel{\vec{\nabla} \phi} \cdot \vec{B} = 0$$



Basics of the model: linear regime

Let's go now deep inside inflation

Dynamical equations:

Inflaton in slow-roll,
Homogeneous

+

dominated by the potential


$$\dot{\tilde{\pi}}_\phi = a \cancel{\vec{\nabla}^2} \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \cancel{\vec{\nabla} \phi} \times \tilde{\vec{E}}$$

- No backreaction in inflation's dynamics
- No generation of inhomogeneities

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \cancel{\vec{\nabla} \phi} \cdot \vec{B} = 0$$



Basics of the model: linear regime

Let's go now deep inside inflation

Dynamical equations:

Inflaton in slow-roll,
Homogeneous

+

dominated by the potential

$$\dot{\tilde{\pi}}_\phi = a \cancel{\vec{\nabla}^2} \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Constraint equation:
Gauss's law

- No backreaction in inflation's dynamics
- No generation of inhomogeneities

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda} \vec{\nabla} \phi \cdot \vec{B} = 0$$

Gauge fields get excited by this term



Basics of the model: linear regime

EoMs

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Constraint equation:
Gauss's law


$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda}\vec{\nabla}\phi \cdot \vec{B} = 0$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$

$$\left(\begin{array}{l} K_\phi \equiv \frac{1}{2}\pi_\phi^2 = \frac{1}{2a^6}\tilde{\pi}_\phi^2, \quad G_\phi \equiv \frac{1}{2a^2} \sum (\partial_i\phi)^2, \quad V \equiv \frac{1}{2}m^2\phi^2, \\ K_A \equiv \frac{1}{2} \sum_i \frac{E_i^2}{a^2} = \frac{1}{2} \sum_i \frac{\tilde{E}_i^2}{a^4}, \quad G_A \equiv \frac{1}{2} \sum_i \frac{B_i^2}{a^4} \end{array} \right)$$



Basics of the model: linear regime

EoMs

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Constraint equation:
Gauss's law


$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda}\vec{\nabla}\phi \cdot \vec{B} = 0$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$

$$\left(\begin{array}{l} K_\phi \equiv \frac{1}{2}\pi_\phi^2 = \frac{1}{2a^6}\tilde{\pi}_\phi^2, \quad G_\phi \equiv \frac{1}{2a^2} \sum (\partial_i\phi)^2, \quad V \equiv \frac{1}{2}m^2\phi^2, \\ K_A \equiv \frac{1}{2} \sum_i \frac{E_i^2}{a^2} = \frac{1}{2} \sum_i \frac{\tilde{E}_i^2}{a^4}, \quad G_A \equiv \frac{1}{2} \sum_i \frac{B_i^2}{a^4} \end{array} \right)$$



Basics of the model: linear regime

EoMs

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Constraint equation:
Gauss's law

$$\vec{\nabla} \cdot \tilde{\vec{E}} + \frac{1}{\Lambda}\vec{\nabla}\phi \cdot \vec{B} = 0$$


Gauge fields evolve in
the background set by
the axion field

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \left\langle -2K_\phi + V + K_A - G_A \right\rangle$$

$$\begin{pmatrix} K_\phi \equiv \frac{1}{2}\pi_\phi^2 = \frac{1}{2a^6}\tilde{\pi}_\phi^2, & G_\phi \equiv \frac{1}{2a^2} \sum (\partial_i\phi)^2, & V \equiv \frac{1}{2}m^2\phi^2, \\ K_A \equiv \frac{1}{2} \sum_i \frac{E_i^2}{a^2} = \frac{1}{2} \sum_i \frac{\tilde{E}_i^2}{a^4}, & G_A \equiv \frac{1}{2} \sum_i \frac{B_i^2}{a^4} \end{pmatrix}$$



Basics of the model: linear regime

An analytical solution can be found


Changing to the helicity basis: + Conformal time

$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i \vec{k} \cdot \vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$



Basics of the model: linear regime

An analytical solution can be found

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$


Changing to the helicity basis: + Conformal time

$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k} \cdot \vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis: + Conformal time


$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states


$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$



$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis: + Conformal time


$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$


$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$



$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis:

+

Conformal time


$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$




$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Two polarisations:

- One exponentially amplified
- The other not

Chiral instability



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis:

+

Conformal time

$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$


$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Two polarisations:

- One exponentially amplified
- The other not

Chiral instability



Basics of the model: linear regime

An analytical solution can be found

Changing to the helicity basis:

+

Conformal time

$$\vec{A}(\vec{x}, \tau) = \sum_{\lambda=\pm} \int \frac{d^3 k}{(2\pi)^3} A^\lambda(k, \tau) \vec{\varepsilon}^\lambda(\hat{k}) e^{i\hat{k}\cdot\vec{x}}$$

Photon with 2 helicity states

$$k_i \varepsilon_i^\pm(\hat{k}) = 0, \quad \epsilon_{ijk} k_j \varepsilon_k^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k})$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}), \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(\hat{k}) = \delta_{\lambda, \lambda'}$$

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$


$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Two polarisations:

- One exponentially amplified
- The other not

Chiral instability

$$|A^+| \gg |A^-|$$




Basics of the model: linear regime

Analytical solution:

[M. M. Anber, L. Sorbo (0908.4089)]

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda} \quad \left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

Instability controlled by ξ



Basics of the model: linear regime

Analytical solution:


[M. M. Anber, L. Sorbo (0908.4089)]

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda} \quad \left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

Instability controlled by ξ

Deep inside inflation $\xi \rightarrow \text{const}$

$$A^+(k, t) = \frac{e^{\frac{\pi}{2}\xi}}{\sqrt{2k}} W_{i\xi, \frac{1}{2}}(2ikt) \simeq \frac{1}{\sqrt{2k}} \left(\frac{k}{2|\xi|aH} \right)^{1/4} e^{\pi|\xi|-2\sqrt{2|\xi|k/aH}}$$



Basics of the model: linear regime

Analytical solution:

[M. M. Anber, L. Sorbo (0908.4089)]

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda} \quad \left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

Instability controlled by

ξ

Deep inside inflation $\xi \rightarrow \text{const}$

$$A^+(k, t) = \frac{e^{\frac{\pi}{2}\xi}}{\sqrt{2k}} W_{i\xi, \frac{1}{2}}(2ikt) \simeq \frac{1}{\sqrt{2k}} \left(\frac{k}{2|\xi|aH} \right)^{1/4} e^{\pi|\xi|-2\sqrt{2|\xi|k/aH}}$$


Topological term:

Appears in EoMs

$$\frac{1}{a^3} \langle \vec{E} \cdot \vec{B} \rangle \simeq 2.4 \cdot 10^{-4} \frac{H^4}{|\xi|^4} e^{2\pi|\xi|}$$

EM energy:

$$\frac{1}{2a^4} \langle a^2 E^2 + B^2 \rangle \simeq 1.4 \cdot 10^{-4} \frac{H^4}{|\xi|^3} e^{2\pi|\xi|}$$



Basics of the model: linear regime

Analytical solution:

[M. M. Anber, L. Sorbo (0908.4089)]

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda} \quad \left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

Instability controlled by

ξ

Deep inside inflation

$\xi \rightarrow \text{const}$

$$A^+(k, t) = \frac{e^{\frac{\pi}{2}\xi}}{\sqrt{2k}} W_{i\xi, \frac{1}{2}}(2ikt) \simeq \frac{1}{\sqrt{2k}} \left(\frac{k}{2|\xi|aH} \right)^{1/4} e^{\pi|\xi|-2\sqrt{2|\xi|k/aH}}$$

Topological term:


Appears in EoMs

$$\frac{1}{a^3} \langle \vec{E} \cdot \vec{B} \rangle \simeq 2.4 \cdot 10^{-4} \frac{H^4}{|\xi|^4} e^{2\pi|\xi|}$$

EM energy:

$$\frac{1}{2a^4} \langle a^2 E^2 + B^2 \rangle \simeq 1.4 \cdot 10^{-4} \frac{H^4}{|\xi|^3} e^{2\pi|\xi|}$$

Amplified around
the Hubble scale
at each time



Basics of the model: linear regime

Analytical solution:

[M. M. Anber, L. Sorbo (0908.4089)]

Topological term:

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

Instability controlled by ξ

Deep inside inflation

$$\xi \rightarrow \text{const}$$

$$A^+(k, t) = \frac{e^{\frac{\pi}{2}\xi}}{\sqrt{2k}} W_{i\xi, \frac{1}{2}}(2ikt) \simeq \frac{1}{\sqrt{2k}} \left(\frac{k}{2|\xi|aH} \right)^{1/4} e^{\pi|\xi|-2\sqrt{2|\xi|k/aH}}$$

ξ

Not accurate towards the end of inflation


Appears in EoMs

$$\frac{1}{a^3} \langle \vec{E} \cdot \vec{B} \rangle \simeq 2.4 \cdot 10^{-4} \frac{H^4}{|\xi|^4} e^{2\pi|\xi|}$$

EM energy:

$$\frac{1}{2a^4} \langle a^2 E^2 + B^2 \rangle \simeq 1.4 \cdot 10^{-4} \frac{H^4}{|\xi|^3} e^{2\pi|\xi|}$$


Amplified around the Hubble scale at each time




Basics of the model: linear regime

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

-6 eFolds
↓
0 eFolds







Basics of the model: linear regime

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

-6 eFolds
↓
0 eFolds






Basics of the model: linear regime

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_{\pm}(k, \tau) = 0$$

-6 eFolds


0 eFolds




Basics of the model: linear regime

[J. R. C. Cuissa, D. G. Figueroa (1812.03132)]

Energy components




Topological term vs slow-roll




Basics of the model: linear regime

[J. R. C. Cuissa, D. G. Figueroa (1812.03132)]


Energy components



Topological term vs slow-roll



Backreaction effect cannot be neglected




Basics of the model: linear regime

Linear regime

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$



Basics of the model: linear regime

Linear regime

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi\vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$

Basics of the model: linear regime

Linear regime


Dynamical equations:

$$\begin{aligned} \dot{\tilde{\pi}}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}} \end{aligned}$$

Neglected

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) = \frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$



Basics of the model

Beyond the linear regime: backreaction


Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi\vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:


$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$
$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

Backreaction terms



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle ,$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$


$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]

[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]

+

[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle ,$$
$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a \Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$


*1st approximation
homogeneous*

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

- [G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]
[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]
+
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

1st approximation
homogeneous

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle ,$$
$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

Expansion equation:


$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$
$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]

[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]

+

[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle,$$
$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$
$$\langle \vec{E} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} |A_+(\vec{k}, \tau)|^2$$

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]


[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]

+

[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$
$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:


$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \tilde{\vec{E}} \cdot \vec{B} \rangle,$$
$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$
$$\langle \tilde{\vec{E}} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} \left| A_+(\vec{k}, \tau) \right|^2$$

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)
[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]

+
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$
$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$
$$\langle K_A + G_A \rangle = \frac{1}{a^4} \int \frac{dk}{4\pi^2} k^2 \left(\left| \frac{dA_+(\vec{k}, \tau)}{d\tau} \right|^2 + k^2 |A_+(\vec{k}, \tau)|^2 \right)^2$$



Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \vec{E} \cdot \vec{B} \rangle,$$
$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$
$$\langle \vec{E} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} |A_+(\vec{k}, \tau)|^2$$

1st approximation
homogeneous
Due to
backreaction
not only
slow-roll

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)
[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]
+
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$
$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$
$$\langle K_A + G_A \rangle = \frac{1}{a^4} \int \frac{dk}{4\pi^2} k^2 \left(\left| \frac{dA_+(\vec{k}, \tau)}{d\tau} \right|^2 + k^2 |A_+(\vec{k}, \tau)|^2 \right)^2$$

Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \vec{E} \cdot \vec{B} \rangle,$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\langle \vec{E} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} |A_+(\vec{k}, \tau)|^2$$

Due to
backreaction
not only
slow-roll

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]
[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]

+
[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

$$\langle K_A + G_A \rangle = \frac{1}{a^4} \int \frac{dk}{4\pi^2} k^2 \left(\left| \frac{dA_+(\vec{k}, \tau)}{d\tau} \right|^2 + k^2 |A_+(\vec{k}, \tau)|^2 \right)^2$$

Basics of the model

Beyond the linear regime: backreaction

Dynamical equations:

*1st approximation
homogeneous*

$$\tilde{\pi}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a \Lambda} \langle \vec{E} \cdot \vec{B} \rangle,$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\langle \vec{E} \cdot \vec{B} \rangle = -\frac{\lambda}{a^4} \int \frac{dk}{4\pi} k^3 \frac{d}{d\tau} |A_+(\vec{k}, \tau)|^2$$

Due to
backreaction
not only
slow-roll

[G. Dall'Agata, S. González-Martín, A. Papageorgiou, M. Peloso (1912.09950)]

[V. Domcke, V. Guidetti, Y. Welling A. Westphal (2002.02952)]


[E. V. Gorbar, K. Schmitz, O. O. Sobol, S. I. Vilchinskii, (2109.01651)]

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} (\langle -2K_\phi + V \rangle - \langle K_A + G_A \rangle)$$

$$\langle K_A + G_A \rangle = \frac{1}{a^4} \int \frac{dk}{4\pi^2} k^2 \left(\left| \frac{dA_+(\vec{k}, \tau)}{d\tau} \right|^2 + k^2 |A_+(\vec{k}, \tau)|^2 \right)^2$$




Why in the lattice?

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi\vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\begin{aligned}\dot{\pi}_a &= \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) = \\ &\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle\end{aligned}$$



Why in the lattice?

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = -a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B},$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}$$

No gradient terms induced
(yet)

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields


Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a\vec{\nabla}^2\phi - a^3m^2\phi + \frac{1}{a\Lambda}\tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi\vec{B} + \frac{1}{a\Lambda}\vec{\nabla}\phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2}(3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2}\langle -2K_\phi + V - K_A - G_A \rangle$$



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

Inhomogeneous source:
Induce gradients not
tractable analytically



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields

Dynamical equations:


$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} - \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

Inhomogeneous source:
Induce gradients not
tractable analytically



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields

Dynamical equations:


$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}\end{aligned}$$

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

Inhomogeneous source:
Induce gradients not
tractable analytically



Why in the lattice?

Lattice provides the solution:
Real time evolution of fields

Dynamical equations:

$$\begin{aligned}\dot{\tilde{\pi}}_\phi &= a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B}, \\ \dot{\tilde{\vec{E}}} &= -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}\end{aligned}$$

Inhomogeneous source:
Induce gradients not
tractable analytically

Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$



$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

\forall D.o.F will be simultaneously
evolved

- Track of all inhomogeneities
- Full backreaction effects included

Discretisation procedure

[D. G. Figueroa & M. Shaposhnikov (1705.09629)]



ϕ @ sites
 A_μ @ links

Want to preserve:

1- Gauge transformations

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

2- Bianchi identities


$$\vec{\nabla} \times \vec{E} = \dot{\vec{B}} \quad \vec{\nabla} \cdot \vec{B} = 0$$

3- Topological nature of the coupling

$$F_{\mu\nu}\tilde{F}^{\mu\nu} = \partial_\mu K^\mu$$

Exact lattice shift symmetry

4- Continuum limit to $\mathcal{O}(dx^2)$



Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]


Continuum

$$\frac{\phi}{\Lambda} \vec{E} \cdot \vec{B}$$

Lattice

$$\sum_i \frac{\phi}{\Lambda} E_i^{(2)} B_i^{(4)}$$

The only one
that satisfies
all conditions



Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]

Continuum

$$\frac{\phi}{\Lambda} \vec{E} \cdot \vec{B}$$


The only one
that satisfies
all conditions


Lattice

$$\sum_i \frac{\phi}{\Lambda} E_i^{(2)} B_i^{(4)}$$

$$E_i^{(2)} \equiv \frac{1}{2}(E_i + E_{i,-i}) \quad @ \quad n + \frac{\hat{0}}{2}$$

$$B_i^{(4)} \equiv \frac{1}{4}(B_i + B_{i,-j} + B_{i,-k} + B_{i,-j-k}) \quad @ \quad n$$





Discretisation procedure


[D. G. Figueroa, M. Shaposhnikov (1705.09629)]

Continuum

$$\frac{\phi}{\Lambda} \vec{E} \cdot \vec{B}$$

$$E_i^{(2)} \equiv \frac{1}{2}(E_i + E_{i,-i}) \quad @ \quad n + \frac{\hat{0}}{2}$$

$$B_i^{(4)} \equiv \frac{1}{4}(B_i + B_{i,-j} + B_{i,-k} + B_{i,-j-k}) \quad @ \quad n$$



Lattice

$$\sum_i \frac{\phi}{\Lambda} E_i^{(2)} B_i^{(4)}$$

Want to preserve:

1- Gauge transformations

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$

2- Bianchi identities


$$\vec{\nabla} \times \vec{E} = \dot{\vec{B}} \quad \vec{\nabla} \cdot \vec{B} = 0$$

3- Topological nature of the coupling

$$F_{\mu\nu} \tilde{F}^{\mu\nu} = \partial_\mu K^\mu$$

Exact lattice shift symmetry

4- Continuum limit to $\mathcal{O}(dx^2)$



Discretisation procedure


[D. G. Figueroa, M. Shaposhnikov (1705.09629)]

Continuum

$$\frac{\phi}{\Lambda} \vec{E} \cdot \vec{B}$$

$$E_i^{(2)} \equiv \frac{1}{2}(E_i + E_{i,-i}) \quad @ \quad n + \frac{\hat{0}}{2}$$

$$B_i^{(4)} \equiv \frac{1}{4}(B_i + B_{i,-j} + B_{i,-k} + B_{i,-j-k}) \quad @ \quad n$$



Lattice

$$\sum_i \frac{\phi}{\Lambda} E_i^{(2)} B_i^{(4)}$$

Want to preserve:

1- Gauge transformations

$$A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$$


2- Bianchi identities

$$\vec{\nabla} \times \vec{E} = \dot{\vec{B}} \quad \vec{\nabla} \cdot \vec{B} = 0$$

3- Topological nature of the coupling

$$F_{\mu\nu} \tilde{F}^{\mu\nu} = \text{Exact lattice shift symmetry}$$


4- Continuum limit to $\mathcal{O}(dx^2)$



Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$




Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} (\tilde{\pi}_\phi)^2 - \frac{1}{2}a_{+\hat{\frac{0}{2}}} \left(\Delta_i^+ \phi_{+\hat{\frac{0}{2}}} \right)^2 - \frac{1}{2}a_{+\hat{\frac{0}{2}}}^3 m^2 \phi_{+\hat{\frac{0}{2}}} \right. \\ & + \frac{1}{2}a_{+\hat{\frac{0}{2}}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \end{aligned}$$




Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} (\tilde{\pi}_\phi)^2 - \frac{1}{2}a_{+\hat{0}} \left(\Delta_i^+ \phi_{+\hat{0}} \right)^2 - \frac{1}{2}a_{+\hat{0}}^3 m^2 \phi_{+\hat{0}} \right. \\ & + \frac{1}{2}a_{+\hat{0}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \end{aligned}$$




Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} (\tilde{\pi}_\phi)^2 - \frac{1}{2}a_{+\hat{0}} \left(\Delta_i^+ \phi_{+\hat{0}} \right)^2 - \frac{1}{2}a_{+\hat{0}}^3 m^2 \phi_{+\hat{0}} \right. \\ & + \frac{1}{2}a_{+\hat{0}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \end{aligned}$$




Discretisation procedure

[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a \left(\vec{\nabla}\phi \right)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} (\tilde{\pi}_\phi)^2 - \frac{1}{2}a_{+\hat{0}} \left(\Delta_i^+ \phi_{+\hat{0}} \right)^2 - \frac{1}{2}a_{+\hat{0}}^3 m^2 \phi_{+\hat{0}} \right. \\ & + \frac{1}{2}a_{+\hat{0}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \end{aligned}$$



Discretisation procedure


[D. G. Figueroa, M. Shaposhnikov (1705.09629)]
[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

$$S = \int d^4x \left[\frac{1}{2}a^3\pi_\phi^2 - \frac{1}{2}a (\vec{\nabla}\phi)^2 - \frac{1}{2}a^3m^2\phi^2 + \frac{1}{2}a \left(\vec{E}^2 - \frac{\vec{B}^2}{a^2} \right) + \frac{\phi}{\Lambda} \vec{E} \cdot \vec{B} \right]$$

Full discretised action:

$$\begin{aligned} S = & \Delta t \Delta x^3 \sum_{t, \vec{n}} \left[\frac{1}{2a^3} \left(\frac{1}{2}a_{+\frac{\hat{0}}{2}} \left(\Delta_i^+ \phi_{+\frac{\hat{0}}{2}} \right)^2 - \frac{1}{2}a_{+\frac{\hat{0}}{2}} m^2 \phi_{+\frac{\hat{0}}{2}} \right. \right. \\ & + \frac{1}{2}a_{+\frac{\hat{0}}{2}} \sum_i \left(\Delta_0^+ A_i - \Delta_i^+ A_0 \right)^2 - \frac{1}{4a} \sum_{i,j} \left(\Delta_i^+ A_j - \Delta_j^+ A_i \right)^2 \\ & \left. \left. + \frac{\phi}{a\Lambda} \sum_i \frac{1}{2} \tilde{E}_i^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \right] \right] \end{aligned}$$

Ready for variational principle




Discretisation procedure

Common kick-drift scheme

Evolution Kernels

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L , \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi , \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$



Discretisation procedure

Common kick-drift scheme


Evolution Kernels

Dynamical equations:

$$\tilde{\pi}_{\phi,+}{}^{\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L , \quad \tilde{E}_{i,+}{}^{\hat{0}} = \tilde{E}_{i,-}{}^{\hat{0}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+}{}^{\frac{\hat{0}}{2}} = \phi_{-}{}^{\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi , \quad A_{i,+}{}^{\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+}{}^{\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+}{}^{\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+}{}^{\frac{\hat{0}}{2}} - a_{+}^3 m^2 \phi_{+}{}^{\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+}^{(2)} \left(B_i^{(4)} + B_{i,+}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+}{}^i B_{i,+}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+}{}^{\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-}{}^{\frac{\hat{0}}{2}} \right\} \end{aligned}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

Dynamical equations:


$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+ \frac{\hat{0}}{2}} = \tilde{E}_{i,- \frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+ \frac{\hat{0}}{2}} = \phi_{- \frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+ \hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+ \frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+ \frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+ \frac{\hat{0}}{2}} - a_{+ \frac{\hat{0}}{2}}^3 m^2 \phi_{+ \frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+ \frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+ \hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+ \frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{- \frac{\hat{0}}{2}} \right\} \end{aligned}$$

Gauss's constraint:

$$\sum_i \Delta_i^- \tilde{E}_{i,+ \frac{\hat{0}}{2}} = -\frac{1}{4\Lambda} \sum_{\pm} \sum_i \left(\Delta_i^\pm \phi_{+ \frac{\hat{0}}{2}} \right) \left(B_i^{(4)} + B_{i,+ \hat{0}}^{(4)} \right)_{\pm i}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

Dynamical equations:


$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+ \frac{\hat{0}}{2}} = \tilde{E}_{i,- \frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+ \frac{\hat{0}}{2}} = \phi_{- \frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+ \hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+ \frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+ \frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+ \frac{\hat{0}}{2}} - a_{+ \frac{\hat{0}}{2}}^3 m^2 \phi_{+ \frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+ \frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+ \hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+ \frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{- \frac{\hat{0}}{2}} \right\} \end{aligned}$$

Gauss's constraint:

$$\sum_i \Delta_i^- \tilde{E}_{i,+ \frac{\hat{0}}{2}} = -\frac{1}{4\Lambda} \sum_{\pm} \sum_i \left(\Delta_i^\pm \phi_{+ \frac{\hat{0}}{2}} \right) \left(B_i^{(4)} + B_{i,+ \hat{0}}^{(4)} \right)_{\pm i}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels


Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$
$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\} \end{aligned}$$

Gauss's constraint:

$$\sum_i \Delta_i^- \tilde{E}_{i,+\frac{\hat{0}}{2}} = -\frac{1}{4\Lambda} \sum_{\pm} \sum_i \left(\Delta_i^\pm \phi_{+\frac{\hat{0}}{2}} \right) \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right)_{\pm i}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$


$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\} \end{aligned}$$

Gauss's constraint:

Alternative approach in:
[A. Caravano, E. Komatsu, K. D. Lozanov, J. Weller (2204.12874)]

$$\sum_i \Delta_i^- \tilde{E}_{i,+\frac{\hat{0}}{2}} = -\frac{1}{4\Lambda} \sum_{\pm} \sum_i \left(\Delta_i^\pm \phi_{+\frac{\hat{0}}{2}} \right) \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right)_{\pm i}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels


$$b_{+\hat{0}} = b + dt\mathcal{K}_a^L$$

Expansion equations:

$$\mathcal{K}_a^L = -\frac{a_{+\hat{0}/2}}{6m_{pl}^2}(\rho_L + 3p_L)_{+\hat{0}/2}$$

Hubble's constraint:

$$\pi_a^2 = \frac{a^2}{3m_{pl}^2}\rho_L$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels

$$b_{+\hat{0}} = b + dt \mathcal{K}_a^L$$

Expansion equations:

$$\mathcal{K}_a^L = -\frac{a_{+\hat{0}/2}}{6m_{pl}^2} (\rho_L + 3p_L)_{+\hat{0}/2}$$

$$(\rho_L + 3p_L)_{+\hat{0}/2} = 2(\bar{K}_\phi^L + \bar{K}_{\phi,+\hat{0}}^L) - 2\bar{V}_{\phi,+\hat{0}/2}^L + 2K_A^L + (G_A^L + G_{A,+\hat{0}}^L)$$


$$\rho_L = \bar{K}_\phi^L + \frac{1}{2}(\bar{G}_{\phi,-\hat{0}/2}^L + \bar{G}_{\phi,+\hat{0}/2}^L) + \frac{1}{2}(\bar{V}_{\phi,-\hat{0}/2}^L + \bar{V}_{\phi,+\hat{0}/2}^L) + \frac{1}{2}(\bar{K}_{A,-\hat{0}/2}^L + \bar{K}_{A,+\hat{0}/2}^L) + \bar{G}_A^L$$

Hubble's constraint:

$$\pi_a^2 = \frac{a^2}{3m_{pl}^2} \rho_L$$

$$\bar{K}_\phi^L = \frac{1}{N^3} \sum_{\vec{n}} \frac{1}{2a^6} \tilde{\pi}_\phi^2, \quad \bar{G}_\phi^L = \frac{1}{N^3} \sum_{\vec{n}} \frac{1}{2} \sum_i (\Delta_i^+ \phi_{+\hat{0}/2})^2, \quad \bar{V}_\phi^L = \frac{1}{N^3} \sum_{\vec{n}} \frac{1}{2} m^2 \phi_{+\hat{0}/2}^2$$

$$\bar{K}_A^L = \frac{1}{N^3} \sum_{\vec{n}} \sum_i \frac{1}{2a_{+\hat{0}/2}^4} \tilde{E}_{i,+\hat{0}/2}^2, \quad \bar{G}_A^L = \frac{1}{N^3} \sum_{\vec{n}} \sum_i \frac{1}{4a^4} (\Delta_i^+ A_j - \Delta_j^+ A_i)^2$$




Discretisation procedure

Evolution Kernels

Dynamical equations:

$$\begin{aligned}\mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+ \frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+ \frac{\hat{0}}{2}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\}\end{aligned}$$



Discretisation procedure


Evolution Kernels

Dynamical equations:

$$\begin{aligned}\mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+i}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\}\end{aligned}$$

Non-symplectic,
Conjugate momenta in kernels!
[Remember 4th lecture & 1st Topical]

$$\dot{\tilde{E}}_i = \mathcal{K}_{i,A}^L \left[a, b, \phi, A_i, \tilde{E}_i \right]$$



Discretisation procedure


Evolution Kernels

Dynamical equations:

$$\begin{aligned}\mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+ \frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+ \frac{\hat{0}}{2}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)}]_{-\frac{\hat{0}}{2}} \right\}\end{aligned}$$

Non-symplectic,
Conjugate momenta in kernels!
[Remember 4th lecture & 1st Topical]

$$\dot{\tilde{E}}_i = \mathcal{K}_{i,A}^L \left[a, b, \phi, A_i, \tilde{E}_i \right]$$




Discretisation procedure

Common kick-drift scheme

Evolution Kernels

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L , \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi , \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$



Discretisation procedure

Common kick-drift scheme


Evolution Kernels

Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$

$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\begin{aligned} \mathcal{K}_\phi^L &= a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right) \\ \mathcal{K}_{i,A}^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\} \end{aligned}$$



Discretisation procedure

Common kick-drift scheme

Evolution Kernels


Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$
$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a^3} \tilde{\pi}_\phi, \quad A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\mathcal{K}_\phi^L = a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi_{+\frac{\hat{0}}{2}} + \frac{1}{2a\Lambda} \sum_i \tilde{E}_{i,+\frac{\hat{0}}{2}}^{(2)} \left(B_i^{(4)} + B_{i,+\hat{0}}^{(4)} \right)$$
$$\mathcal{K}_{i,A}^L = -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ + \frac{1}{8a\Lambda} (2 + dx \Delta_i^+) \sum_{\pm} \sum_{j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\}$$

Implicit scheme required

We need the value of the gauge's canonical momentum @ 0/2 for obtaining its value @ 0/2!



Discretisation procedure

Common kick-drift scheme

Evolution Kernels


Dynamical equations:

$$\tilde{\pi}_{\phi,+\hat{0}} = \tilde{\pi}_\phi + dt \mathcal{K}_\phi^L, \quad \tilde{E}_{i,+\frac{\hat{0}}{2}} = \tilde{E}_{i,-\frac{\hat{0}}{2}} + dt \mathcal{K}_{i,A}^L$$
$$\phi_{+\frac{\hat{0}}{2}} = \phi_{-\frac{\hat{0}}{2}} + \frac{dt}{a} A_{i,+\hat{0}}$$
$$A_{i,+\hat{0}} = A_i + \frac{dt}{a} \tilde{E}_{i,+\frac{\hat{0}}{2}}$$

$$\mathcal{K}_\phi^L = a_{+\frac{\hat{0}}{2}} \sum_i \Delta_i^- \Delta_i^+ \phi_{+\frac{\hat{0}}{2}} - a_{+\frac{\hat{0}}{2}}^3 m^2 \phi$$
$$\mathcal{K}_{i,A}^L = -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- + \frac{1}{8a} \sum_{\pm j,k} \left\{ \epsilon_{ijk} [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{+\frac{\hat{0}}{2}} + [(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}]_{-\frac{\hat{0}}{2}} \right\}$$


Implicit scheme required

We need the value of the gauge's canonical momentum @ 0/2 for obtaining its value @ 0/2!



ANDER URIO


Our Current Work



Explicit-in-time integrator

Joanes

Ander




Explicit-in-time integrator

Joanes

Ander

If momenta appear in kernels



Explicit-in-time integrator


Joanes

If momenta appear in kernels



Ander

Not solvable by explicit
symplectic integrators



Explicit-in-time integrator

Joanes


Ander

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV




Explicit-in-time integrator

Joanes

Ander


If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?



Explicit-in-time integrator

Joanes

If momenta appear in kernels




Ander

Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?

$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV


Why?

Ander

Solution



$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Ander

Non-symplectic integrators

Not solvable by explicit
symplectic integrators


i.e. LF or VV

Why?

Solution



$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?

Ander

Non-symplectic integrators




Solution



Runge-Kutta



$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?

Ander

Non-symplectic integrators




Solution



Runge-Kutta



$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$



Explicit-in-time integrator

Joanes

If momenta appear in kernels



Not solvable by explicit
symplectic integrators

i.e. LF or VV

Why?

$$\pi_{+\frac{\hat{0}}{2}} = \pi_{-\frac{\hat{0}}{2}} + \mathcal{K}[\pi_{+\frac{\hat{0}}{2}}]dt$$

Solution



Ander


Non-symplectic integrators



(explicit version)

Runge-Kutta


No longer!



Explicit-in-time integrator

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$




Explicit-in-time integrator

RK2 & 2 stages

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$






Explicit-in-time integrator

RK2 & 2 stages

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{aligned} x(t_{n+1}) &= x(t_n) + dt (b_1 G_1 + b_2 G_2) \end{aligned} \right\}$$




Explicit-in-time integrator

RK2 & 2 stages

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$



Explicit-in-time integrator


RK2 & 2 stages

1st order ODE

$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$

$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$



Explicit-in-time integrator

RK2 & 2 stages


1st order ODE

$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$

$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$

$$\text{Constraints} \rightarrow b_1 + b_2 = 1 , \ b_2 c_2 = 1/2 , \ b_2 a_{21} = 1/2$$



Explicit-in-time integrator

RK2 & 2 stages

1st order ODE


$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$

$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$

$$\text{Constraints} \rightarrow b_1 + b_2 = 1 , \quad b_2 c_2 = 1/2 , \quad b_2 a_{21} = 1/2$$

Modified Euler



Explicit-in-time integrator

RK2 & 2 stages

1st order ODE


$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$


$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$

$$\text{Constraints} \rightarrow b_1 + b_2 = 1 , \ b_2 c_2 = 1/2 , \ b_2 a_{21} = 1/2$$

Modified Euler



$$b_1 = 1/2 , \ b_2 = 1/2 , \ c_2 = 1 , \ a_{21} = 1$$



Explicit-in-time integrator

RK2 & 2 stages

1st order ODE


$$\frac{dx}{dt} = v(t, x)$$

$$\left. \begin{array}{l} x(t_{n+1}) = x(t_n) + dt (b_1 G_1 + b_2 G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + c_2 dt, x(t_n) + a_{21} G_1 dt) \end{array} \right\}$$

$$x^{\text{Cont}}(t_{n+1}) - x^{\text{RK2}}(t_{n+1}) = \mathcal{O}(dt^3)$$


$$\text{Constraints} \rightarrow b_1 + b_2 = 1 , \ b_2 c_2 = 1/2 , \ b_2 a_{21} = 1/2$$

Modified Euler




$$b_1 = 1/2 , \ b_2 = 1/2 , \ c_2 = 1 , \ a_{21} = 1$$

$$\boxed{\begin{array}{l} x(t_{n+1}) = x(t_n) + \frac{dt}{2} (G_1 + G_2) \\ G_1 = v(t_n, x(t_n)) \\ G_2 = v(t_n + dt, x(t_n) + G_1 dt) \end{array}}$$




Explicit-in-time integrator



Explicit-in-time integrator

2nd order ODE

$$\frac{d^2x}{dt^2} = \mathcal{K}\left(t, x, \frac{dx}{dt}\right)$$




Explicit-in-time integrator

2nd order ODE

$$\frac{d^2x}{dt^2} = \mathcal{K}\left(t, x, \frac{dx}{dt}\right)$$

1st order ODE RK2 ?



Explicit-in-time integrator


2nd order ODE

$$\frac{d^2x}{dt^2} = \mathcal{K}\left(t, x, \frac{dx}{dt}\right)$$





Coupled 1st order ODEs

1st order ODE RK2 ?





Explicit-in-time integrator







Explicit-in-time integrator






Explicit-in-time integrator





Explicit-in-time integrator


Axion-Inflation using RK2



Explicit-in-time integrator


Axion-Inflation using RK2


$$t_n \equiv (1)$$



Explicit-in-time integrator

Axion-Inflation using RK2


$$t_n \equiv (1) \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n) , \\ A_i^{(1)} = A_i(t_n) , \\ a^{(1)} = a(t_n) , \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n) , \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n) , \\ \pi_a^{(1)} = \pi_a(t_n) , \end{array} \right.$$




Explicit-in-time integrator

Axion-Inflation using RK2


$$t_n \equiv (1) \quad \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n) , \\ A_i^{(1)} = A_i(t_n) , \\ a^{(1)} = a(t_n) , \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n) , \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n) , \\ \pi_a^{(1)} = \pi_a(t_n) , \end{array} \right. \quad \xrightarrow{\text{dotted arrow}} \quad \left\{ \begin{array}{l} K_\phi^{(1)} = K_\phi(a^{(1)}, \tilde{\pi}_\phi^{(1)}) \\ V^{(1)} = V(\phi^{(1)}) \\ K_A^{(1)} = K_A(a^{(1)}, \tilde{E}_i^{(1)}) \\ G_A^{(1)} = G_A(a^{(1)}, A_i^{(1)}) \end{array} \right.$$



Explicit-in-time integrator


Axion-Inflation using RK2


$$\begin{aligned} t_n \equiv (1) & \quad \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n), \\ A_i^{(1)} = A_i(t_n), \\ a^{(1)} = a(t_n), \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n), \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n), \\ \pi_a^{(1)} = \pi_a(t_n), \end{array} \right. \\ & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(1)} = K_\phi(a^{(1)}, \tilde{\pi}_\phi^{(1)}) \\ V^{(1)} = V(\phi^{(1)}) \\ K_A^{(1)} = K_A(a^{(1)}, \tilde{E}_i^{(1)}) \\ G_A^{(1)} = G_A(a^{(1)}, A_i^{(1)}) \end{array} \right. \\ & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(1)} = \mathcal{K}_\phi \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_A^{(1)} = \mathcal{K}_A \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_a^{(1)} = \mathcal{K}_a \left[a^{(1)}, K_\phi^{(1)}, V^{(1)}, K_A^{(1)}, G_A^{(1)} \right]. \end{array} \right. \end{aligned}$$



Explicit-in-time integrator


Axion-Inflation using RK2






Explicit-in-time integrator


Axion-Inflation using RK2






Explicit-in-time integrator


Axion-Inflation using RK2






Explicit-in-time integrator

Axion-Inflation using RK2






Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}
 t_n \equiv (1) & \quad \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n), \\ A_i^{(1)} = A_i(t_n), \\ a^{(1)} = a(t_n), \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n), \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n), \\ \pi_a^{(1)} = \pi_a(t_n), \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(1)} = K_\phi(a^{(1)}, \tilde{\pi}_\phi^{(1)}) \\ V^{(1)} = V(\phi^{(1)}) \\ K_A^{(1)} = K_A(a^{(1)}, \tilde{E}_i^{(1)}) \\ G_A^{(1)} = G_A(a^{(1)}, A_i^{(1)}) \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(1)} = \mathcal{K}_\phi \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_A^{(1)} = \mathcal{K}_A \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_a^{(1)} = \mathcal{K}_a \left[a^{(1)}, K_\phi^{(1)}, V^{(1)}, K_A^{(1)}, G_A^{(1)} \right]. \end{array} \right.
 \end{aligned}$$

$$\begin{aligned}
 t_{n+1} \equiv (2) & \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.
 \end{aligned}$$




Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}
 t_n \equiv (1) & \quad \left\{ \begin{array}{l} \phi^{(1)} = \phi(t_n), \\ A_i^{(1)} = A_i(t_n), \\ a^{(1)} = a(t_n), \\ \tilde{\pi}_\phi^{(1)} = \tilde{\pi}_\phi(t_n), \\ \tilde{E}_i^{(1)} = \tilde{E}_i(t_n), \\ \pi_a^{(1)} = \pi_a(t_n), \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(1)} = K_\phi(a^{(1)}, \tilde{\pi}_\phi^{(1)}) \\ V^{(1)} = V(\phi^{(1)}) \\ K_A^{(1)} = K_A(a^{(1)}, \tilde{E}_i^{(1)}) \\ G_A^{(1)} = G_A(a^{(1)}, A_i^{(1)}) \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(1)} = \mathcal{K}_\phi \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_A^{(1)} = \mathcal{K}_A \left[\phi^{(1)}, \tilde{\pi}_\phi^{(1)}, a^{(1)}, \tilde{E}^{(1)}, \vec{B}^{(1)} \right], \\ \mathcal{K}_a^{(1)} = \mathcal{K}_a \left[a^{(1)}, K_\phi^{(1)}, V^{(1)}, K_A^{(1)}, G_A^{(1)} \right]. \end{array} \right.
 \end{aligned}$$


$$\begin{aligned}
 t_{n+1} \equiv (2) & \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \\
 & \xrightarrow{\text{dotted arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.
 \end{aligned}$$



Explicit-in-time integrator

Axion-Inflation using RK2


$$\begin{aligned} t_{n+1} \equiv (2) & \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t , \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t , \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t , \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t , \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t , \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t . \end{array} \right. \\ & \quad \xrightarrow{\text{dotted blue arrow}} \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \\ & \quad \xrightarrow{\text{dotted blue arrow}} \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right] , \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right] , \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right] . \end{array} \right. \end{aligned}$$



Explicit-in-time integrator

Axion-Inflation using RK2

$$t_{n+1} \equiv (2) \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \quad \xrightarrow{\text{dotted blue arrows}} \quad \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \quad \xrightarrow{\text{dotted blue arrow}} \quad \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.$$



$$\phi(t_{n+1}) = \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2$$


$$A_i(t_{n+1}) = A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2$$

$$a(t_{n+1}) = a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2$$

$$\tilde{\pi}_\phi(t_{n+1}) = \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t$$

$$\tilde{E}_i(t_{n+1}) = \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t$$


$$\pi_a(t_{n+1}) = \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t$$



Explicit-in-time integrator

Axion-Inflation using RK2

$$t_{n+1} \equiv (2) \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \quad \xrightarrow{\text{dotted blue arrows}} \quad \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \quad \xrightarrow{\text{dotted blue arrow}} \quad \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.$$



$$\phi(t_{n+1}) = \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2$$


$$A_i(t_{n+1}) = A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2$$

$$a(t_{n+1}) = a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2$$

$$\tilde{\pi}_\phi(t_{n+1}) = \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t$$

$$\tilde{E}_i(t_{n+1}) = \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t$$


$$\pi_a(t_{n+1}) = \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t$$



Explicit-in-time integrator

Axion-Inflation using RK2

$$t_{n+1} \equiv (2) \quad \left\{ \begin{array}{l} \phi^{(2)} = \phi^{(1)} + (a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} \delta t, \\ A_i^{(2)} = A_i^{(1)} + (a^{(1)})^{-1} \tilde{E}_i^{(1)} \delta t, \\ a^{(2)} = a^{(1)} + \pi_a^{(1)} \delta t, \\ \tilde{\pi}_\phi^{(2)} = \tilde{\pi}_\phi^{(1)} + \mathcal{K}_\phi^{(1)} \delta t, \\ \tilde{E}_i^{(2)} = \tilde{E}_i^{(1)} + \mathcal{K}_A^{(1)} \delta t, \\ \pi_a^{(2)} = \pi_a^{(1)} + \mathcal{K}_a^{(1)} \delta t. \end{array} \right. \quad \xrightarrow{\text{dotted blue arrows}} \quad \left\{ \begin{array}{l} K_\phi^{(2)} = K_\phi(a^{(2)}, \tilde{\pi}_\phi^{(2)}) \\ V^{(2)} = V(\phi^{(2)}) \\ K_A^{(2)} = K_A(a^{(2)}, \tilde{E}_i^{(2)}) \\ G_A^{(2)} = G_A(a^{(2)}, A_i^{(2)}) \end{array} \right. \quad \xrightarrow{\text{dotted blue arrow}} \quad \left\{ \begin{array}{l} \mathcal{K}_\phi^{(2)} = \mathcal{K}_\phi \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_A^{(2)} = \mathcal{K}_A \left[\phi^{(2)}, \tilde{\pi}_\phi^{(2)}, a^{(2)}, \tilde{E}_i^{(2)}, \vec{B}^{(2)} \right], \\ \mathcal{K}_a^{(2)} = \mathcal{K}_a \left[a^{(2)}, K_\phi^{(2)}, V^{(2)}, K_A^{(2)}, G_A^{(2)} \right]. \end{array} \right.$$



$$\phi(t_{n+1}) = \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2$$


$$A_i(t_{n+1}) = A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2$$

$$a(t_{n+1}) = a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2$$

$$\tilde{\pi}_\phi(t_{n+1}) = \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t$$

$$\tilde{E}_i(t_{n+1}) = \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t$$

$$\pi_a(t_{n+1}) = \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t$$




Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$




Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$

More intermediate
kernels





Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$






Explicit-in-time integrator


Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$



More intermediate
kernels





Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$







Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$

$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$







Explicit-in-time integrator

Axion-Inflation using RK2

$$\begin{aligned}\phi(t_{n+1}) &= \phi(t_n) + \left[(a^{(1)})^{-3} \tilde{\pi}_\phi^{(1)} + (a^{(2)})^{-3} \tilde{\pi}_\phi^{(2)} \right] \delta t / 2 \\ A_i(t_{n+1}) &= A_i(t_n) + \left[(a^{(1)})^{-1} \tilde{E}_i^{(1)} + (a^{(2)})^{-1} \tilde{E}_i^{(2)} \right] \delta t / 2 \\ a(t_{n+1}) &= a(t_n) + (\pi_a^{(1)} + \pi_a^{(2)}) \delta t / 2\end{aligned}$$


$$\begin{aligned}\tilde{\pi}_\phi(t_{n+1}) &= \tilde{\pi}_\phi(t_n) + \frac{1}{2} (\mathcal{K}_\phi^{(1)} + \mathcal{K}_\phi^{(2)}) \delta t \\ \tilde{E}_i(t_{n+1}) &= \tilde{E}_i(t_n) + \frac{1}{2} (\mathcal{K}_A^{(1)} + \mathcal{K}_A^{(2)}) \delta t \\ \pi_a(t_{n+1}) &= \pi_a(t_n) + \frac{1}{2} (\mathcal{K}_a^{(1)} + \mathcal{K}_a^{(2)}) \delta t\end{aligned}$$





Explicit-in-time integrator

Axion-Inflation using RK2



Explicit-in-time integrator

Axion-Inflation using RK2


Dynamical equations

$$\mathcal{K}_\phi^L = a \sum_i \Delta_i^- \Delta_i^+ \phi - a^3 m^2 \phi + \frac{1}{2a\Lambda} \sum_i \tilde{E}_i^{(2)} B_i^{(4)}$$

$$\mathcal{K}_A^L = -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right)$$

$$+ \frac{1}{4a\Lambda} \sum_{\pm} \sum_{j,k} \epsilon_{ijk} \left\{ \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_{+i} + \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right] \right\}$$

$$\mathcal{K}_a^L = -\frac{a}{6m_{pl}^2} (\rho_L + 3p_L)$$



Explicit-in-time integrator


Axion-Inflation using RK2

Dynamical equations

$$\begin{aligned}\mathcal{K}_\phi^L &= a \sum_i \Delta_i^- \Delta_i^+ \phi - a^3 m^2 \phi + \frac{1}{2a\Lambda} \sum_i \tilde{E}_i^{(2)} B_i^{(4)} \\ \mathcal{K}_A^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{4a\Lambda} \sum_{\pm} \sum_{j,k} \epsilon_{ijk} \left\{ \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_{+i} + \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_i \right\} \\ \mathcal{K}_a^L &= -\frac{a}{6m_{pl}^2} (\rho_L + 3p_L)\end{aligned}$$

Constraints

$$\begin{aligned}\sum_i \Delta_i^- \tilde{E}_i &= -\frac{1}{2\Lambda} \sum_{\pm} \sum_i (\Delta_i^\pm \phi) B_{i,\pm i}^{(4)} \\ \pi_a^2 &= \frac{a^2}{3m_{pl}^2} \rho_L\end{aligned}$$



Explicit-in-time integrator

Axion-Inflation using RK2

No temporal semi-sums

Dynamical equations

$$\mathcal{K}_\phi^L = a \sum_i \Delta_i^- \Delta_i^+ \phi - a^3 m^2 \phi + \frac{1}{2a\Lambda} \sum_i \tilde{E}_i^{(2)} B_i^{(4)}$$

$$\mathcal{K}_A^L = -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right)$$


$$+ \frac{1}{4a\Lambda} \sum_{\pm} \sum_{j,k} \epsilon_{ijk} \left\{ \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_{+i} + \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_i \right\}$$

$$\mathcal{K}_a^L = -\frac{a}{6m_{pl}^2} (\rho_L + 3p_L)$$

Constraints

$$\sum_i \Delta_i^- \tilde{E}_i = -\frac{1}{2\Lambda} \sum_{\pm} \sum_i (\Delta_i^\pm \phi) B_{i,\pm i}^{(4)}$$

$$\pi_a^2 = \frac{a^2}{3m_{pl}^2} \rho_L$$



Explicit-in-time integrator


Axion-Inflation using RK2

Dynamical equations

$$\begin{aligned}\mathcal{K}_\phi^L &= a \sum_i \Delta_i^- \Delta_i^+ \phi - a^3 m^2 \phi + \frac{1}{2a\Lambda} \sum_i \tilde{E}_i^{(2)} B_i^{(4)} \\ \mathcal{K}_A^L &= -\frac{1}{a} \sum_{j,k} \epsilon_{ijk} \Delta_j^- B_k - \frac{1}{2a^3 \Lambda} \left(\tilde{\pi}_\phi B_i^{(4)} + \tilde{\pi}_{\phi,+i} B_{i,+i}^{(4)} \right) \\ &\quad + \frac{1}{4a\Lambda} \sum_{\pm} \sum_{j,k} \epsilon_{ijk} \left\{ \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_{+i} + \left[(\Delta_j^\pm \phi) \tilde{E}_{k,\pm j}^{(2)} \right]_i \right\} \\ \mathcal{K}_a^L &= -\frac{a}{6m_{pl}^2} (\rho_L + 3p_L)\end{aligned}$$

Constraints


$$\begin{aligned}\sum_i \Delta_i^- \tilde{E}_i &= -\frac{1}{2\Lambda} \sum_{\pm} \sum_i (\Delta_i^\pm \phi) B_{i,\pm i}^{(4)} \\ \pi_a^2 &= \frac{a^2}{3m_{pl}^2} \rho_L\end{aligned}$$




Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme

[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]







Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme


[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]






Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme





Explicit-in-time integrator


Axion-Inflation using RK2: validation of the scheme


[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

Spectrum of A

$$\frac{k^3}{2\pi^2 m^2} |A_k|^2$$

Preheating example





Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme

[J. R. C. Cuissa , D. G. Figueroa (1812.03132)]

Spectrum of A

$$\frac{k^3}{2\pi^2 m^2} |A_k|^2$$

Preheating example


Comoving momentum


$$k/m$$

Solid: Canivete and Figueroa
Scatter: Our scheme

LF implicit

RK2 explicit







Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme

[J. R. C. Cuissa , D. .G. Figueroa (1812.03132)]







Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme

[J. R. C. Cuissa , D. .G. Figueroa (1812.03132)]







Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme


[J. R. C. Cuissa , D. .G. Figueroa (1812.03132)]






Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme







Explicit-in-time integrator

Axion-Inflation using RK2: validation of the scheme


Gauss constraint
observable





Initial Conditions in the Linear Regime


$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$



Initial Conditions in the Linear Regime

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis → Conformal time




Initial Conditions in the Linear Regime

$$\dot{\tilde{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$

Helicity basis → Conformal time

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$


Helicity basis



Conformal time

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis


Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$



$$|A^+| \gg |A^-|$$



Initial Conditions in the Linear Regime


$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$



$$|A^+| \gg |A^-|$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$



Initial Conditions in the Linear Regime


$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis → Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$


$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Deep inside Hubble $k \gg aH$



$$|A^+| \gg |A^-|$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis → Conformal time


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

Deep inside Hubble $k \gg aH$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0 \quad \left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$

$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$

$|A^+| \gg |A^-|$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis



Conformal time


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution


Deep inside Hubble $k \gg aH$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$



$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$

$$|A^+| \gg |A^-|$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis



Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$


Bunch-Davies solution

Deep inside Hubble $k \gg aH$


$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$




$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$



$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$



$|A^+| \gg |A^-|$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis



Conformal time


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution


Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$



$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$



$|A^+| \gg |A^-|$

Cosmic time

$\tau \simeq -1/aH$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a}\vec{\nabla} \times \vec{B} - \frac{1}{a^3\Lambda}\tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution

Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$


$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$

$$\left(\frac{\partial^2}{\partial\tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$


Cosmic time


$$\tau \simeq -1/aH$$

$$A_+(k, t) \simeq \frac{1}{\sqrt{2k}} e^{ik/a(t)H(t)}$$



Initial Conditions in the Linear Regime





Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution

Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$


Cosmic time

$\tau \simeq -1/aH$

$$A_+(k, t) \simeq \frac{1}{\sqrt{2k}} e^{ik/a(t)H(t)}$$

Time derivative

$$E_+(k, t) \simeq \frac{i}{a} \sqrt{\frac{k}{2}} e^{ik/a(t)H(t)}$$



Initial Conditions in the Linear Regime

$$\dot{\vec{E}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B}$$

Helicity basis

Conformal time

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \pm \frac{2k\xi}{\tau} \right) A_\pm(k, \tau) = 0$$

$$\xi \equiv \frac{\dot{\phi}}{2H\Lambda}$$

Bunch-Davies solution

Deep inside Hubble $k \gg aH$

$$A_+(k, \tau) \simeq \frac{1}{\sqrt{2k}} e^{-ik\tau}$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 \right) A_+(k, \tau) \simeq 0$$

$$\left(\frac{\partial^2}{\partial \tau^2} + k^2 + \frac{2k\xi}{\tau} \right) A_+(k, \tau) = 0$$


Cosmic time

$\tau \simeq -1/aH$

$$A_\pm(k, t) \simeq \frac{1}{\sqrt{2k}} e^{ik/a(t)H(t)}$$

Time derivative


$$E_\pm(k, t) \simeq \frac{i}{a} \sqrt{\frac{k}{2}} e^{ik/a(t)H(t)}$$



Initial Conditions in the Linear Regime


Helicity basis

BD solutions for A^\pm, E^\pm





Initial Conditions in the Linear Regime







Initial Conditions in the Linear Regime






Initial Conditions in the Linear Regime






Initial Conditions in the Linear Regime




$$k_i \varepsilon_i^\pm(\hat{k}) = 0 , \quad \epsilon_{ijk} k_j \varepsilon_i^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k}) ,$$

$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}) , \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(-\hat{k}) = \delta_{\lambda\lambda'}$$




Initial Conditions in the Linear Regime



$$k_i \varepsilon_i^\pm(\hat{k}) = 0 , \quad \epsilon_{ijk} k_j \varepsilon_i^\pm(\hat{k}) = \mp i k \varepsilon_i^\pm(\hat{k}) ,$$


$$\varepsilon_i^\pm(\hat{k})^* = \varepsilon_i^\pm(-\hat{k}) , \quad \varepsilon_i^\lambda(\hat{k}) \varepsilon_i^{\lambda'}(-\hat{k}) = \delta_{\lambda\lambda'}$$

$$A_i^\pm = P_{ij}^\pm A_j$$



Initial Conditions in the Linear Regime

How to build the
Helicity Projector?




Initial Conditions in the Linear Regime

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?



Initial Conditions in the Linear Regime


Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



Initial Conditions in the Linear Regime

How to build the
Helicity Projector?


Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

2 eigenvalues: 1 and -1

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



Initial Conditions in the Linear Regime


Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



Initial Conditions in the Linear Regime

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?




Helicity projector

$$P_{ij}^{\pm} = \frac{1}{2} (\Sigma_{ij}^{\pm} + (\Sigma^{\pm})_{ij}^2)$$

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



Initial Conditions in the Linear Regime

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?




Helicity projector


$$P_{ij}^{\pm} = \frac{1}{2} (\Sigma_{ij}^{\pm} + (\Sigma^{\pm})_{ij}^2)$$

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



$$P_{ij}^{\pm} = \frac{1}{2} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \pm \frac{i}{k} \epsilon_{ijk} k_k \right)$$



Initial Conditions in the Linear Regime

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$

How to build the
Helicity Projector?




Helicity projector

$$P_{ij}^{\pm} = \frac{1}{2} (\Sigma_{ij}^{\pm} + (\Sigma^{\pm})_{ij}^2)$$

Transverse projector


$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$



$$P_{ij}^{\pm} = \frac{1}{2} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \pm \frac{i}{k} \epsilon_{ijk} k_k \right)$$



$$A_i^{\pm} = P_{ij}^{\pm} A_j$$
$$E_i^{\pm} = P_{ij}^{\pm} E_j$$



Initial Conditions in the Linear Regime

How to build the
Helicity Projector?

Helicity operator

$$\Sigma_{ij}^{\pm}(k) = \pm \frac{i}{k} \epsilon_{ijk} k_k$$




Helicity projector

$$P_{ij}^{\pm} = \frac{1}{2} (\Sigma_{ij}^{\pm} + (\Sigma^{\pm})_{ij}^2)$$

Transverse projector

$$(\Sigma^{\pm})_{ij}^2(k) = \delta_{ij} - \frac{k_i k_j}{k^2}$$




$$P_{ij}^{\pm} = \frac{1}{2} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \pm \frac{i}{k} \epsilon_{ijk} k_k \right)$$



$$A_i^{\pm} = P_{ij}^{\pm} A_j$$
$$E_i^{\pm} = P_{ij}^{\pm} E_j$$

Gauss


$$\vec{\nabla} \cdot \vec{E} = 0$$



Initial Conditions in the Linear Regime

Two IC options






Initial Conditions in the Linear Regime

Two IC options

One option

Only project the + helicity states






Initial Conditions in the Linear Regime

Two IC options

One option

Only project the + helicity states

Why?



Initial Conditions in the Linear Regime


Two IC options

One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced



Initial Conditions in the Linear Regime

Two IC options


One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced

Loss of information!



Initial Conditions in the Linear Regime

One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced




Two IC options

Our option

Project both states + and -





Initial Conditions in the Linear Regime


Two IC options

One option

Only project the + helicity states

Why?


A^+ is exponentially enhanced



Our option

Project both states + and -

Why?



Initial Conditions in the Linear Regime


Two IC options

One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced




Our option

Project both states + and -

Why?

Main objective: non-linear regime





Initial Conditions in the Linear Regime


One option

Only project the + helicity states

Why?

A^+ is exponentially enhanced

Loss of information!



Two IC options

Our option


Project both states + and -

Why?

Main objective: non-linear regime




Can A^- become relevant?



Initial Conditions in the Linear Regime

Gauge initialization process




Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$




Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF



Initial Conditions in the Linear Regime

Gauge initialization process


- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

- 2 Add fluctuations to A_i with RGF



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables


$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

- 2 Add fluctuations to A_i with RGF

- 4 Repeat the process with A_-



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$


- 2 Add fluctuations to A_i with RGF

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

- 4 Repeat the process with A_-

Fourier



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF


- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

Fourier

- 4 Repeat the process with A_-

- 5 Go back to real space via FFT



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$


Fourier

- 4 Repeat the process with A_-

- 5 Go back to real space via FFT



EOMs evolved in real space!



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$


Fourier

- 4 Repeat the process with A_-

- 5 Go back to real space via FFT

- 6 Repeat initialization with E^\pm BD solutions

EOMs evolved in real space!



Initial Conditions in the Linear Regime

Gauge initialization process

- 1 Assign BD + solution to A_i CL variables

$$A_i \rightarrow \frac{1}{\sqrt{2k}} e^{ik/aH}$$

- 2 Add fluctuations to A_i with RGF

- 3 Use the helicity projector

$$A_i \rightarrow A^+ = P_{ij}^+ A_j$$

Fourier

- 4 Repeat the process with A_-


- 5 Go back to real space via FFT


- 6 Repeat initialization with E^\pm BD solutions

EOMs evolved in real space!


⋮ We will have an amplitude of 2 BD in A_i
$$\sum_i |A_i|^2 = |A_+|^2 + |A_-|^2$$


Simulations in the Linear Regime







Simulations in the Linear Regime







Simulations in the Linear Regime







Simulations in the Linear Regime





Simulations in the Linear Regime






Non-linear evolution: example


Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$

$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$

@





Non-linear evolution: example

Dynamical equations:

$$\dot{\tilde{\pi}}_\phi = a \vec{\nabla}^2 \phi - a^3 m^2 \phi + \frac{1}{a\Lambda} \tilde{\vec{E}} \cdot \vec{B},$$


$$\dot{\tilde{\vec{E}}} = -\frac{1}{a} \vec{\nabla} \times \vec{B} - \frac{1}{a^3 \Lambda} \tilde{\pi}_\phi \vec{B} + \frac{1}{a\Lambda} \vec{\nabla} \phi \times \tilde{\vec{E}}$$


Expansion equation:

$$\dot{\pi}_a = \frac{-a}{6m_{\text{pl}}^2} (3p + \rho) =$$

$$\frac{a}{3m_{\text{pl}}^2} \langle -2K_\phi + V - K_A - G_A \rangle$$

@







Non-linear evolution: example

Gauge PS

A^+

Linear







Non-linear evolution: example

Gauge PS

A^+

Linear






Non-linear evolution: example


Gauge PS


A^+

Linear



Non-linear






Non-linear evolution: example


Gauge PS


A^+

Linear



Non-linear







Non-linear evolution: example

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}$$

Linear







Non-linear evolution: example

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}$$

Linear







Non-linear evolution: example

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}$$

Linear

Non-linear







Non-linear evolution: example

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}$$

Linear


Non-linear






Non-linear evolution: example


What's
happening in
those extra
folds?






Non-linear evolution: example

What's
happening in
those extra
folds?







Non-linear evolution: example

What's
happening in
those extra
folds?

A^+
 A_{tot}






Non-linear evolution: example

What's
happening in
those extra
folds?

A^+

A_{tot}

A^-





Non-linear evolution: example

Inflation observables

Linear

Non-linear







7- Non-linear evolution: example

Inflation observables

Linear

Non-linear







7- Non-linear evolution: example

Inflation observables

Linear

Non-linear







7- Non-linear evolution: example

Inflation observables

Linear

Non-linear






Summary & Conclusions

- Successful implementation of axion-inflation in CL


$$\frac{\phi}{\Lambda} F \tilde{F}$$



Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simpletic integrators needed: RK
 - Validation: preheating

$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$




Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simpletic integrators needed: RK
 - Validation: preheating




$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$



Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simplectic integrators needed: RK
 - Validation: preheating
- Initial conditions
 - Helicity basis
 - Validation: linear regime


$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$



Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simplectic integrators needed: RK
 - Validation: preheating
- Initial conditions
 - Helicity basis
 - Validation: linear regime


$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$



Summary & Conclusions


- Successful implementation of axion-inflation in CL
- Non-simplectic integrators needed: RK
 - Validation: preheating
- Initial conditions
 - Helicity basis
 - Validation: linear regime
- Preliminari non-linear sims already show inhomogeneities non-negligible

$$\boxed{\frac{\phi}{\Lambda} F \tilde{F}}$$



Summary & Conclusions

- Successful implementation of axion-inflation in CL
- Non-simplectic integrators needed: RK
 - Validation: preheating
- Initial conditions
 - Helicity basis
 - Validation: linear regime
- Preliminari non-linear sims already show inhomogeneities non-negligible



Ready to study non-linear phenomenology!

$$\frac{\phi}{\Lambda} F \tilde{F}$$

Thank you!